



## Original Article

## The Epworth Sleepiness Scale's Completeness Difficulties in an Urban Cameroonian Population

### *Difficultés à répondre à l'échelle de somnolence d'Epworth dans une population camerounaise urbaine*

Massongo Massongo<sup>1</sup>, Ekono Bitchong CF<sup>2,3</sup>, Bassogbag R<sup>4</sup>, Afane Ze E<sup>1,2</sup>

#### RÉSUMÉ

**Introduction.** In order to check some empirical observations, we aimed to assess difficulties in answering the Epworth Sleepiness Scale (ESS) and estimate its completeness rate in urban young, healthy and literate population in Cameroon. **Methodology.** This cross-sectional study was conducted on 2016, among randomly selected adults in Yaoundé, the capital of Cameroon. We collected sociodemographic and occupational data, as well as the ESS. The later assessed the sleepiness risk in 8 different daily situations : Sitting and reading, watching television, sitting inactive in a public place, passenger in a car for an hour without a break, lying down to rest in the afternoon, talking to someone, sitting quietly after an alcohol-free lunch and being in a car while stopped for a few minutes in the traffic. Unanswered items were noted, their number and distribution were described. **Results.** A total of 400 Yaoundé inhabitants (mean age  $34.8 \pm 11.1$  years, 57% men, 60.8% office workers) were enrolled. One participant (0.25%) could not answer the item 1, 3 (0.75%) could not answer the item 4 and 237 (59.2%) were limited with the item 8. Overall, 239 participants could not complete the ESS, giving a completeness rate [95% confident interval] of 40.3 [35.5; 45.1]. Among them, 236 were let down by a single item, 2 by 2 items and 1 by 3. **Interpretation.** This study confirmed difficulties in completing the ESS, even in a young and literate population. Further research is needed to confirm these difficulties with the ESS completeness, especially in rural populations and other regions. **Conclusion.** Epworth sleepiness scale is not easy to complete, even for a young, healthy and literate Cameroonian population.

#### ABSTRACT

**Introduction.** Afin de vérifier certaines observations empiriques, nous avons cherché à évaluer les difficultés à répondre à l'échelle de somnolence d'Epworth (ESS) et à estimer son taux d'exhaustivité dans la population urbaine jeune, saine et alphabétisée du Cameroun. **Méthodologie.** Cette étude transversale a été réalisée en 2016, auprès d'adultes sélectionnés au hasard à Yaoundé, la capitale du Cameroun. Nous avons collecté des données socio-démographiques et professionnelles, ainsi que l'ESS. Ce dernier a évalué le risque de somnolence dans 8 situations quotidiennes différentes: s'asseoir et lire, regarder la télévision, rester inactif dans un lieu public, passager dans une voiture pendant une heure sans interruption, allongé pour se reposer l'après-midi, parler à quelqu'un, assis tranquillement après un déjeuner sans alcool et être dans une voiture pendant quelques minutes dans la circulation. Les éléments restés sans réponse ont été notés, leur nombre et leur distribution ont été décrits. **Résultats.** Au total, 400 habitants de Yaoundé (âge moyen de  $34,8 \pm 11,1$  ans, 57% d'hommes, 60,8% d'employés de bureau) étaient inscrits. Un participant (0,25%) n'a pas pu répondre au point 1, 3 (0,75%) n'ont pas pu répondre au point 4 et 237 (59,2%) ont été limités au point 8. Dans l'ensemble, 239 participants n'ont pas pu compléter l'ESS, donnant un taux d'exhaustivité [intervalle de confiance à 95%] de 40,3 [35,5; 45,1]. Parmi eux, 236 ont été déçus par un seul élément, 2 par 2 éléments et 1 par 3. **Interprétation.** Cette étude a confirmé les difficultés à terminer l'ESS, même dans une population jeune et alphabétisée. Des recherches supplémentaires sont nécessaires pour confirmer ces difficultés avec l'exhaustivité du ESS, en particulier dans les populations rurales et d'autres régions. **Conclusion.** L'échelle de somnolence d'Epworth n'est pas facile à compléter, même pour une population camerounaise jeune, en bonne santé et alphabétisée.

<sup>1</sup>. Faculty of Medicine and Biomedical Sciences, University of Yaoundé 1, Yaoundé, Cameroon

<sup>2</sup>. Jamot Hospital of Yaoundé, Yaoundé, Cameroon

<sup>3</sup>. Faculty of Medicine and Pharmaceutical Sciences, University of Douala, Douala, Cameroon

<sup>4</sup>. Higher Institute of Medical Technology, Nkolondom, Cameroon

#### Auteur correspondant :

Massongo Massongo,  
Department of Medicine and Specialties, Faculty of Medicine and Biomedical Sciences, University of Yaoundé 1, Yaoundé, Cameroon

Adresse e-mail :

[massongom@gmail.com](mailto:massongom@gmail.com)

**Mots-clés :** Epworth sleepiness scale, Completeness, difficulties, Sub-Saharan Africa.

#### Keywords:

Échelle de somnolence d'Epworth, Complétude, difficultés, Afrique subsaharienne.

#### INTRODUCTION

Chronic non communicable diseases (NCD) have a growing prevalence in sub-Saharan Africa (SSA). This

includes diabetes, high blood pressure, dyslipidemia, obesity and cardiovascular diseases [1]–[7]. Sleep related

disorders (SRD), including the obstructive sleep apnea syndrome (OSA), are part of these NCD. Data on OSA are still scarce in SSA. However, recent community studies revealed OSA's prevalence (5 – 10%) close to the one of the western world [8]–[11]. Furthermore, Benjafield published in 2019 a modelling-based meta-analysis, estimating OSA prevalence in almost all countries of the world. This prevalence ranged 7.8% – 77% for an apnea-hypopnea index (AHI)  $\geq$  5/h and 3 – 36.6% for moderate to severe OSA (AHI  $\geq$  15/h). African countries' prevalence estimates were not lower than others. Cameroon ones were 36.6% and 14.7% for global OSA and moderate to severe OSA respectively [12]. Excessive daytime sleepiness (EDS) is a major symptom in sleep related disorders and OSA especially. It is also associated with cardiovascular diseases, quality of life and other health problems [13]–[19]. The objective diagnosis of SRD and EDS requires sophisticated procedures such as ventilatory polygraphy, polysomnography, iterative sleep latency tests and maintenance of wakefulness test. Access to those procedures are still limited, especially in resource-limited settings as SSA, where screening tools are of great importance. The Epworth sleepiness scale (ESS) is one of the most used screening tool for EDS [20] [20]. However, some items of the ESS are not designed to be universally answered (see methodology section). Our daily practice revealed frequent difficulties to complete the form. Those difficulties have been assumed to be related to socio economic status or other factors such as rural area, access to electricity, literacy or access to public transport. We aimed to check these observations by estimating the completeness rate of the ESS in urban young, healthy and literate population in Cameroon.

## MATERIAL AND MÉTHODS

### *Study design, settings and population*

This cross-sectional study took place in Yaoundé, the political capital of Cameroon, between November 2015 and May 2016. Black subjects aged 16 or over and living in the city and its surroundings were asked to participate. We selected the population study using a two-stage clusters stratified sampling. The stratification variable was the type of activity, divided into office activity (administrative buildings) and outdoor activity (neighborhoods and / or markets). In the office activity group: 2 buildings in the city (Ministry of Higher Education and National Institute of Youth and Sport) were first chosen by convenience, then one or more services from each institution were randomly chosen, and all the members of the selected services were systematically invited to participate to the study, until the expected number of participants was obtained. In the outdoor activity group: 4 districts (Omnisport, Essos, Tropicana and Ekounou) were chosen by convenience among the most popular and endowed with significant external activity (markets, street vendors, manual workers). Then the streets of each district were crossed, and the questionnaire was proposed to all working adults encountered.

### *Data collection and management*

Data were collected during a face-to-face interview. Seventh year medical students specifically trained for the purpose conducted the interviews. The questionnaire included socio-demographic features (age, gender, place of residence, marital status), lifestyle and habits (occupation, toxics and drugs consumptions, activity profile using the Ricci and Gagnon auto test), personal and family medical history (particularly cardiovascular and metabolic conditions), symptoms enquiry, physical examination, and Epworth sleepiness scale (ESS). The ESS evaluates the sleepiness risk in 8 different daily situations, relating to usual way of life in recent times, which are :

1. Sitting and reading,
2. Watching television,
3. Sitting inactive in a public place (e.g. a theatre or a meeting),
4. As a passenger in a car for an hour without a break,
5. Lying down to rest in the afternoon when circumstances permit
6. Sitting and talking to someone
7. Sitting quietly after a lunch without alcohol
8. In a car, while stopped for a few minutes in the traffic

Each item is marked from 0 (would never doze or fall asleep) to 3 (high chance of dozing or falling asleep), giving a total score range of 0 to 24. Excessive daytime sleepiness is defined by an ESS  $>$  10 [20]. ESS was administered as an auto-questionnaire, in the presence of the investigator who could explain questions and items to participants, but not suggest answers.

The ESS completeness rate was defined as the proportion of respondents who answered to the 8 items without difficulties. Continuous variables were expressed as mean  $\pm$  standard deviation, and categorical ones as counts and percentages. Epidata version 3.1 was used to digitize data, and Stata version 12.0 was used for data analysis.

### *Ethical statements*

The study was approved by the Institutional Ethics Committee for Research on Human Health, at the University of Douala. A written consent was given by each participant before its enrollment.

## RÉSULTATS

### *Baseline characteristics of study participants*

A total of 400 subjects were enrolled in our study, including 228 men (sex ratio = 1.3). They had a mean age of  $34.8 \pm 11.2$  years. Two-fifths (41.3%) of them were aged 25 to 34 years and the extreme ages were 16 and 86. The place of residence was exclusively urban (69.5%) or peri-urban (30.5%). The occupation was considered as physically inactive (office worker or no occupation) for 62.8% of the sample. Alcohol consumption and known HBP were present in nearly half and a third of them respectively, while almost 40% were obese or overweight. A little proportion of respondents confessed illegal drugs consumption (See Table 1).

**Table 1: Baseline characteristics of subjects enrolled in the study on the Epworth sleepiness scale completeness rate**

Characteristic	Modality	Values
Age, years	Mean +/- standard deviation	34,8 ± 11,1
	Extremes	16 - 86
Sex, number	men/women	228/172
Place of residence	Urban	278 (69.5%)
	Peri-urban	122 (30.5%)
Occupation type	Manual worker	130 (32.5%)
	Itinerant worker	19 (4.7%)
	Office worker	243 (60.8%)
	No activity	8 (2.0%)
Activity score (Nina and Gagnon)	Inactive	18 (4.5%)
	Active	375 (93.7%)
	Very active	7 (1.8%)
History of high blood pressure		138 (34.5%)
Alcohol consumption		196 (49.0%)
Tobacco consumption		21 (5.2%)
Coffee consumption		142 (35.5%)
Illegal drug consumption		7 (1.7%)
Psychotropic medication		21 (5.2%)
Weight category (N=310)	Thin	12 (3.9%)
	Normal weight	178 (57.4%)
	Overweight	80 (25.8%)

### ESS completeness and difficulties

Two hundred and thirty nine out of the 400 participants (59.7%) could not. Thus, the ESS completeness rate [95% confident interval] was 40.3 [35.5 ; 45.1].

Item 8 (In a car, while stopped for a few minutes in the traffic) was the most involved in failure to complete ESS, with 237 (59.2%) persons concerned. The 2 other uncompleted items were 4 (As a passenger in a car for an hour without a break) and 1 (Sitting and reading), with 3 and 1 participants unable to respond respectively (See figure 1).




Figure 1 : Number of participants unable to answer properly for each item, in the study on the Epworth sleepiness scale completeness rate

Regarding the number of items, 236 participants (98.7%) out of the 239 who were unable to complete the ESS had a difficulty for a single item, while 2 were concerned by 2 items and the last one by 3 items (see figure 2).




Figure 2 : Number of participants unable to respond properly, with respect to the number of items concerned, in the study on the Epworth sleepiness scale completeness rate.

### DISCUSSION

In this cross-sectional study of the ESS answering difficulties, more than half of participants could not complete the questionnaire. Most of them were limited by a single item, which appeared to be the 8<sup>th</sup> one (In a car, while stopped for a few minutes in the traffic).

In our knowledge, there are no published data on the ESS completeness rate in sub-Saharan Africa, and this issue is poorly documented even in other regions, especially on young and healthy populations. A study conducted among 104 nondemented older subjects revealed that 60% of them could not answer at least one question on the ESS [21]. Another one found a proportion of 64% in a geriatric Caucasian population with a mean age of 82 ± 8 years [22]. One could assume that the elderly is not the suitable population for a rigorous assessment of the ESS, since cognitive or sensorial defects (known or not) may alter their abilities to respond. However, this highlights the fact that our data were alarming, as our participants were young, healthy and active, and we found comparable results.

In a Brazilian study designed to compare sleepiness scores in 231 adults with different levels of blood pressure, 60 of them (26%) were excluded due to difficulties in the obtained data, including the ESS. This study was the sole we found that addressed difficulties with ESS in adult (no elderly) population. But not reporting or publishing targeted data on this issue does not mean that they don't exist. On the ESS form, respondents are asked to imagine themselves in unknown situations as follows: "Even if you haven't done some of these things recently, try to work out how they would have affected you" [20]. But this may be quite a difficult exercise, increasing risk of wrong answers. It has been shown that ESS scores were significantly different when the questionnaires were filled by the patients themselves rather than their relatives [23] or their caregivers [24].

In this resource-limited setting, having the item 8 overrepresented was not surprising, since few people own a car and/or have ever driven one. Very few people were limited by more than 1 item. In our opinion, the number and proportions of items concerned would be quite greater if the study included rural areas. Items such

as reading, watching television or being a passenger in a car for an hour, are expected to be more impacted in rural areas. This assumption is supported by the following facts : 1) 23% of adults Cameroonian are illiterate and most of illiterate live in rural areas [25], 2) 39% of the population don't have access to electricity in Cameroon, and the situation is worse in rural areas [26], 3) The population/car ratio in Cameroon is as low as 15 cars/1 000 inhabitants [27], and 4) public transportation means in rural areas usually do not allow any sleepiness (cars overloading, bad roads, use of bikes as taxi and bike overloading, etc.).

One strength of our methodology was the absence of intervention from our investigator in the responses' formulation. This permitted the real expression of potential difficulties. Enrolling only in urban and peri-urban population was a limitation, since this was not representative of the whole country which have more than 60% of rural population [28]. However, this was a voluntary strategy intended to reveal ESS limitations in the less unfavored people.

This study confirmed difficulties in completing the ESS, even in a young and literate population. Further research is needed to confirm these difficulties with the ESS completeness, especially in rural populations and in the elderly, as it has been shown elsewhere. Physicians and other caregivers should accompany patients while filling the questionnaire, in order to minimize the potential disruption of the results. However, clinicians would welcome a new EDS screening tool, which would have less limitations related to socio-economic aspects or personal possessions.

## CONCLUSION

Epworth sleepiness scale is not easy to complete, even for a young, healthy and literate Cameroonian population.

## RÉFÉRENCES

1. L. K. Fezeu *et al.*, « Ten-year changes in central obesity and BMI in rural and urban Cameroon », *Obes. Silver Spring Md*, vol. 16, n° 5, p. 1144-1147, mai 2008, doi: 10.1038/oby.2008.44.
2. L. Fezeu, A. P. Kengne, B. Balkau, P. K. Awah, et J. C. Mbanya, « Ten-year change in blood pressure levels and prevalence of hypertension in urban and rural Cameroon », *J. Epidemiol. Community Health*, vol. 64, n° 4, p. 360-365, avr. 2010, doi: 10.1136/jech.2008.086355.
3. A. P. Kengne, P. K. Awah, L. Fezeu, et J. C. Mbanya, « The burden of high blood pressure and related risk factors in urban sub-Saharan Africa: evidences from Douala in Cameroon », *Afr. Health Sci.*, vol. 7, n° 1, p. 38-44, mars 2007, doi: 10.5555/afhs.2007.7.1.38.
4. P. Kengne et P. K. Awah, « Classical cardiovascular risk factors and all-cause mortality in rural Cameroon », *QJM Mon. J. Assoc. Physicians*, vol. 102, n° 3, p. 209-215, mars 2009, doi: 10.1093/qjmed/hcn175.
5. J. B. Echouffo-Tcheugui et A. P. Kengne, « Chronic non-communicable diseases in Cameroon - burden, determinants and current policies », *Glob. Health*, vol. 7, p. 44, nov. 2011, doi: 10.1186/1744-8603-7-44.
6. F. Verdier et L. Fourcade, « Changes in cardiovascular risk factors in developing countries », *Med. Trop. Rev. Corps Sante Colon.*, vol. 67, n° 6, p. 552-558, déc. 2007.
7. R. M. Kamadjeu, R. Edwards, J. S. Atanga, E. C. Kiawi, N. Unwin, et J.-C. Mbanya, « Anthropometry measures and prevalence of obesity in the urban adult population of Cameroon: an update from the Cameroon Burden of Diabetes Baseline Survey », *BMC Public Health*, vol. 6, p. 228, sept. 2006, doi: 10.1186/1471-2458-6-228.
8. Kenne Kenyo, « Prévalence et déterminants du syndrome d'apnées-hypopnées obstructives du sommeil à Bandjoun », Thèse de doctorat en médecine, Institut Supérieur de Technologie Médicale, Yaoundé, 2016.
9. O. B. Ozoh *et al.*, « Prospective assessment of the risk of obstructive sleep apnea in patients attending a tertiary health facility in Sub-Saharan Africa », *Pan Afr. Med. J.*, vol. 17, p. 302, 2014, doi: 10.11604/pamj.2014.17.302.2898.
10. M. Massongo, A. D. Balkissou, V. Poka, S. Voufouo Sonwa, et E. W. Pefura Yone, « Syndrome d'apnées de sommeil et comorbidités cardiovasculaire, métabolique et psychiatrique : une étude hospitalière », *EM-Consulte*. <https://www.em-consulte.com/article/1198658/article/syndrome-d-apnees-de-sommeil-et-comorbidites-cardi> (consulté le juin 19, 2019).
11. N. C. Netzer *et al.*, « Prevalence of symptoms and risk of sleep apnea in primary care », *Chest*, vol. 124, n° 4, p. 1406-1414, oct. 2003.
12. A. V. Benjafield *et al.*, « Estimation of the global prevalence and burden of obstructive sleep apnoea: a literature-based analysis », *Lancet Respir. Med.*, vol. 7, n° 8, p. 687-698, août 2019, doi: 10.1016/S2213-2600(19)30198-5.
13. J. Xie *et al.*, « Excessive Daytime Sleepiness Independently Predicts Increased Cardiovascular Risk After Myocardial Infarction », *J. Am. Heart Assoc.*, vol. 7, n° 2, janv. 2018, doi: 10.1161/JAHA.117.007221.
14. P. E. Ronksley *et al.*, « Excessive daytime sleepiness is associated with increased health care utilization among patients referred for assessment of OSA », *Sleep*, vol. 34, n° 3, p. 363-370, mars 2011, doi: 10.1093/sleep/34.3.363.
15. A. C. Hayley, L. J. Williams, G. A. Kennedy, M. Berk, S. L. Brennan, et J. A. Pasco, « Prevalence of excessive daytime sleepiness in a sample of the Australian adult population », *Sleep Med.*, vol. 15, n° 3, p. 348-354, mars 2014, doi: 10.1016/j.sleep.2013.11.783.
16. Y.-T. Lee, C.-F. Tsai, W.-C. Ouyang, A. C. Yang, C.-H. Yang, et J.-P. Hwang, « Daytime sleepiness: a risk factor for poor social engagement among the elderly », *Psychogeriatr. Off. J. Jpn. Psychogeriatr. Soc.*, vol. 13, n° 4, p. 213-220, déc. 2013, doi: 10.1111/psyg.12020.

17. M. Partinen et C. Guilleminault, « Daytime sleepiness and vascular morbidity at seven-year follow-up in obstructive sleep apnea patients », *Chest*, vol. 97, n° 1, p. 27-32, janv. 1990.
18. E. C. A. Pereira *et al.*, « Prevalence of excessive daytime sleepiness and associated factors in women aged 35-49 years from the "Pindamonhangaba Health Project" (PROSAPIN) », *Rev. Assoc. Médica Bras.* 1992, vol. 58, n° 4, p. 447-452, août 2012.
19. S. Wu, R. Wang, X. Ma, Y. Zhao, X. Yan, et J. He, « Excessive daytime sleepiness assessed by the Epworth Sleepiness Scale and its association with health related quality of life: a population-based study in China », *BMC Public Health*, vol. 12, p. 849, 2012, doi: 10.1186/1471-2458-12-849.
20. M. W. Johns, « A new method for measuring daytime sleepiness: the Epworth sleepiness scale », *Sleep*, vol. 14, n° 6, p. 540-545, déc. 1991, doi: 10.1093/sleep/14.6.540.
21. F. Onen, T. Moreau, N. S. Gooneratne, C. Petit, B. Falissard, et S. H. Onen, « Limits of the Epworth Sleepiness Scale in older adults », *Sleep Breath. Schlaf Atm.*, vol. 17, n° 1, p. 343-350, mars 2013, doi: 10.1007/s11325-012-0700-8.
22. H. Frohnhofen, R. Popp, V. Willmann, H. C. Heuer, et A. Firat, « Feasibility of the Epworth Sleepiness Scale in a sample of geriatric in-hospital patients », *J. Physiol. Pharmacol. Off. J. Pol. Physiol. Soc.*, vol. 60 Suppl 5, p. 45-49, nov. 2009.
23. H. Kumru, J. Santamaria, et R. Belcher, « Variability in the Epworth sleepiness scale score between the patient and the partner », *Sleep Med.*, vol. 5, n° 4, p. 369-371, juill. 2004, doi: 10.1016/j.sleep.2004.03.003.
24. M. Kaminska, V. Jobin, P. Mayer, R. Amyot, M. Perraton-Brillon, et F. Bellemare, « The Epworth Sleepiness Scale: self-administration versus administration by the physician, and validation of a French version », *Can. Respir. J.*, vol. 17, n° 2, p. e27-34, avr. 2010, doi: 10.1155/2010/438676.
25. Banque mondiale, « Taux d'alphabétisation, total des adultes (% des personnes âgées de 15 ans et plus) | Data ». <https://donnees.banquemondiale.org/indicateur/SE.ADT.LITR.ZS?view=chart> (consulté le avr. 12, 2020).
26. Banque mondiale, « Accès à l'électricité (% de la population) | Data ». <https://donnees.banquemondiale.org/indicateur/EG.ELC.ACCS.ZS?view=chart> (consulté le avr. 12, 2020).
27. « List of countries by vehicles per capita », *Wikipedia*. avr. 07, 2020, Consulté le: avr. 12, 2020. [En ligne]. Disponible sur: [https://en.wikipedia.org/w/index.php?title=List\\_of\\_countries\\_by\\_vehicles\\_per\\_capita&oldid=949687704](https://en.wikipedia.org/w/index.php?title=List_of_countries_by_vehicles_per_capita&oldid=949687704).
28. Banque mondiale, « Population urbaine | Data ». <https://donnees.banquemondiale.org/indicateur/SP.URB.TOTL?view=chart> (consulté le avr. 12, 2020).