



Article Original

Prevalence and Factors Associated with Cardiovascular Emergencies in the Emergency Department of Niamey National Hospital

Prévalence et facteurs associés des urgences cardiovasculaires au service des urgences de l'Hôpital National de Niamey

Mounkaila F^{1,2}, Kabore J^{3*}, Yanogo PK^{1,4}, Laurent M^{1,2}, Otshudiandjeka JB^{1,5}, Meda N^{1,4}

Emergency. Prevalence.

¹. Burkina Field Epidemiology and Laboratory Training Program (BFELTP) University Joseph KI-ZERBO, Ouagadougou, Burkina Faso

². Ministry of Health, Niger

³. National Center for Scientific and Technological Research, Institute for Research in Health Sciences, Ouagadougou, Burkina Faso

⁴. Faculty of Medicine, Joseph KI ZERBO University, Ouagadougou, Burkina Faso

⁵. African Field Epidemiology Network (AFENET)

* Corresponding author:

Jean KABORE
jck6238@gmail.com

Keywords: Emergency, Prevalence, cardiovascular, Niger, odds ratio, risk factors

Mots-clés : Urgence, Prévalence, cardiovasculaire, Niger, rapport de cotes, facteurs de risque

ABSTRACT

Introduction. Although many conditions cardiovascular diseases present as emergencies., no study was conducted in Niger to estimate their prevalence and identify related-factors. **Methods.** We conducted a cross-sectional study from February 1st to March 25th, 2020. We included 422 patients aged 15 years or more from the Emergency Department of the National Niamey Hospital (NNH) except those with uncomplete medical records. We carried out questionnaire-based interviews with patients and we collected data from medical files. Data were processed with Epi Info® 7.2.2.6 and MS Excel® 2019. Dependent variable was cardiovascular emergency regardless of the type. We performed a multiple logistic regression model to estimate adjusted odds ratios (95% Confident interval) of the relation between cardiovascular emergencies and patients' characteristics with a statistical significance threshold of p-value < 0.05. **Results.** Globally, 27.5% (116/422) of patients were admitted for CVD emergencies. Among them, 68% (79/116) were over 50 years and 50.9% (59) were male. The most common personal medical history was hypertension with 50.8% (59/116). Among the 15 deaths which were recorded, one third was due to stroke, the main cause of CVD emergency (32/116, 32%). Both hypertensive crisis and heart failure concerned 46.5% (54/116) of all emergencies. History of hypertension [6,09 (3,31-11,21), p <0.0001] and age over than 50 (2.34 [1,33 – 4.10], p=0.002) were independently associated with CVD emergencies. **Conclusion.** Health promoting strategies targeting adequate management of high blood pressure and positive lifestyle habits in people over 50 years could help in reducing cardiovascular emergencies frequency in hospital settings.

RÉSUMÉ

Introduction. De nombreuses pathologies cardiovasculaires se présentent sous la forme d'urgences. Toutefois, aucune étude n'a été menée au Niger pour estimer leur prévalence et identifier les facteurs connexes. **Méthodologie.** Nous avons mené une étude transversale du 1er février au 25 mars 2020. Nous avons inclus 422 patients de ≥15 ans du service d'urgence médicale de l'hôpital national de Niamey (NNH), à l'exception de ceux dont le dossier médical n'était pas complet. Nous avons mené des entretiens par questionnaire avec des patients et nous avons recueilli des données à partir de dossiers médicaux. Les données ont été traitées avec les logiciels Epi Info® 7.2.2.6 et MS Excel® 2019. La variable dépendante était l'urgence cardiovasculaire quel que soit le type. Nous avons effectué un modèle de régression logistique multiple pour estimer les rapports de cotes ajustés (intervalle de confiance à 95 %) de la relation entre les urgences cardiovasculaires et les caractéristiques des patients avec un seuil de signification statistique de valeur p < 0,05. **Résultats.** Au total, 27,5 % (116/422) des patients ont été admis pour des urgences cardiovasculaires. Parmi eux, 68 % (79/116) avaient plus de 50 ans et 50,9 % (59) étaient des hommes. Les antécédents médicaux personnels les plus courants étaient l'hypertension artérielle avec 50,8% (59/116). Parmi les 15 décès enregistrés, un tiers était dû à un accident vasculaire cérébral, principale cause d'urgence cardiovasculaire (32/116, 32%). Les poussées hypertensives et l'insuffisance cardiaque concernaient 46,5% (54/116) de toutes les urgences. Les antécédents d'hypertension [6,09 (3,31-11,21), p <0,0001] et l'âge de plus de 50 ans (2,34 [1,33 – 4,10], p = 0,002) étaient indépendamment associés aux urgences cardiovasculaires. **Conclusion.** Des stratégies de promotion de la santé ciblant une prise en charge adéquate de l'hypertension artérielle et des habitudes de vie positives chez les personnes de plus de 50 ans pourraient aider à réduire la fréquence des urgences cardiovasculaires en milieu hospitalier.

INTRODUCTION

Non-communicable diseases are a set of pathologies with common epidemiological characteristics and

cardiovascular diseases are the leading cause of disease burden in the world (refer to Roth Ag aet al, 2020, in

Kidney int). According to World Health Organization(WHO), 71% of deaths worldwide are related to non-communicable diseases (NCDs), which makes them a major public health issue [1].

Deaths from cardiovascular disease are estimated at 17.7 million, or 31% of total global mortality. Of these deaths 7.4 million are due to coronary heart disease and 6.7 million to stroke (2015) [2]. Main risk factors for coronary heart disease and stroke are poor diet, lack of physical activity, smoking and excessive alcohol use [2]. Many of these cardiovascular conditions occur as emergencies which threaten the patient's immediate life-threatening prognosis. Cardiological emergencies are a set of pathologies such coronary, hemodynamic, rhythmic or of any other origin etc, involving the patient's vital and functional prognosis.

Prevalence of cardiovascular diseases is increasing in sub-Saharan Africa (SSA) [3] [4] and they are the second cause of death in Africa according to WHO' estimates. In 2015, nearly 1.2 million people are dead because of cardiovascular disease, which is more than pooled deaths from malaria and tuberculosis [5]. The main pathologies are high blood pressure, strokes, cardiomyopathies and coronary heart disease.

A study suggests high prevalence of arterial hypertension in children and adolescents in Africa, overweight and obesity being an important risk factor [6]. Also, other authors suggest an increasing burden of stroke in Africa [7].

In Niger, health situation is strongly marked by the predominance of communicable diseases, NCDs are gaining ground. Indeed, according to the "STEPS wise" survey (2007) on risk factors for NCDs and the prevalence of high blood pressure and hyperglycemia, 21.2 % of adults have high blood pressure and 4.3% hyperglycemia. Furthermore, 26.5% of participants were overweight and 4.9% were tobacco smokers [8]. Since then, no national study has been carried out. With a view to curbing the increase in these pathologies, the National Program for Fight against NCDs was created in 2012 and it is leading the design, the implementation and the evaluation of the national policy in fight against NCDs in Niger. Niamey National hospital (NNH) medical emergency receives all medical emergencies and in 2018, about 2,699 cases of cardiovascular emergencies (CVE) were registered.

In our knowledge, there is no study on this issue despite the fact that 21% of admissions to medical emergencies were cardiovascular emergencies (national hospital report). We aim to estimate the prevalence and to identify risk factors of CVE in patients admitted to the medical emergency department of NNH.

METHOD

Type, period and study population

This was a descriptive and analytical cross-sectional study running from February 01 to March 25, 2020. The population consisted of all patients aged 15 years and over, admitted to medical emergency department of the Niamey National Hospital.

Sample size and sampling method

The sample size was calculated considering a proportion of 50%, a precision d of 5% with a statistical power of 80%. Globally, 422 patients were included in the study. We had exhaustively identified all patients aged 15 and over, admitted for a cardiovascular emergency or not, during our study period until reaching our sample size.

Data collection

We conducted a face-to-face interview with patient or his companions if he is not able to respond. A documentary review was also used based on the medical file or patient file in order to complete information on the management. We used a semi-structured questionnaire. These tools were used after sensitizing and obtaining informed consent from patients, in order to collect patient information. We used also medical emergency hospitalization register to identify admitted patients as well as individual medical records to assess information from the preliminary medical examination. we collected sociodemographic (age, sex, occupation, Physical activity, type of physical activity, lifestyle, Socioeconomic factors, Psychological and socio-cultural factors, Place of residence, Marital status, Type of household, Monthly income) and clinical data (Known cardiovascular risk factors, Blood pressure on admission, Height, Weight, body mass index (BMI) abdominal perimeter, Glasgow scale, Diagnosis retained, Outcome).

Statistical analyses

Dependent variable was cardiovascular emergency and independent variables were socio-demographic factors, clinical and paraclinical characteristics, lifestyle, psychological and socio-cultural factors, as well as management. Anthropometric parameters (weight, height, waist circumference) and blood pressure were also measured. With Epi Info[®] version 7.2.2.6 and the Excel[®] version 2019 software's, we estimated frequencies, proportions, means and made comparisons (chi-square tests of Pearson or Student), crude and adjusted odds ratios (OR) with their confidence interval of 95%. The significance level was a p-value <0.05 for all our analyses. We selected variables to be included in the multiple logistic regression model by considering the associations between explanatory variables and the dependent variable with a p-value ≤ 0.20 . Also, we included other variables in the model based on data from scientific literature on the role of these factors in cardiovascular emergencies. We used a top-down walkthrough to identify independently associated factors.

Ethical considerations

This study was conducted within strict framework of compliance with the principles of scientific research. Prior to the administration of the questionnaire, the informed consent of the respondents was required. Also, the study protocol was submitted to the BFELTP' coordination staff for validation. Based on questionnaire, we collected the first and last names of the respondents and then anonymized them. The database was secured in a computer with access conditional on entering a password held by the study investigator.

RESULTS

Prevalence of CVE at the Niamey National Hospital

Among 422 patients included in our study, 116 had cardiovascular emergencies, prevalence of 27.5%)

The most frequent cardiovascular emergencies were stroke, hypertensive crisis, and global heart failure respectively 32% (37), 28% (33), and 18% (21).




Figure 1: Distribution of cardiovascular emergencies at the Niamey National Hospital (February to March 2020)

Distribution of deaths

Among the 15 deaths recorded, 33.3% (5) were due to ischemic stroke.




Figure 2: Distribution of deaths by cardiovascular emergencies at the National Hospital of Niamey from February 1st to March 25th, 2020

Patient data according to CVE status

The mean age was 57.7 (16.7) with extremes of 15 and 90 years. Sixty-eight percent (79) of patients were over 50 years old and approximately half 50.9% (59) were male. Among patients with CVE, 57.7 % (67) did not practice any physical activity, 16.3% (54) were obese and 40.6% (43) had a waist circumference over gender-related standards (for men, a waist circumference ≥ 102 cm and for women a waist circumference ≥ 88 cm). We noted 29.3% (123) had blood pressure $\geq 140/90$ mm Hg. The most common medical history was hypertension (50.8% (59)).

Table I: Comparison of patient data according to cardiovascular emergencies status, Niamey National Hospital, February to March 2020

Data	CVE* (n = 116) %(n)	No CVE (n=306) %(n)	p-value
Age, years			$<10^{-4}$
<50	32 (37)	56 (172)	
≥ 50	68 (79)	44 (134)	
Sex			0.38
Male	51 (59)	55.6 (170)	
Female	49 (57)	44.4 (136)	
Personal history of hypertension			$<10^{-4}$
No	49 (57)	86 (262)	
Yes	51 (59)	14 (44)	
Body mass index, kg/m²			0.7
< 30	86 (100)	87.58 (268)	
≥ 30	14 (16)	12.41 (38)	
Practice of a physical activity			0.01
No	58 (67)	70 (214)	
Yes	42 (49)	30 (92)	
Daily time spent sitting			0.13
<6h	45 (52)	53 (162)	
$\geq 6h$	55 (64)	47 (144)	
Waist circumference[†], n=312			0.02
\leq standard / sex	59.4 (63)	72 (148)	
> standard / sex	40.6 (43)	28 (58)	
Smoking			0.19
No	93 (108)	89 (272)	
Yes	7 (8)	11 (34)	
Spouse smoking, n = 298			$<10^{-4}$
No	96 (111)	99 (296)	
Yes	4 (5)	1 (2)	
Diabetes			0.13
No	94 (109)	89 (273)	
Yes	6 (7)	11 (33)	

*CVE: Cardiovascular emergencies; [†]Waist circumference: for men, we considered a threshold below the standards as a waist circumference of <102 cm and < 88 for women cm.

Crude and adjusted association

In our study, having a history of hypertension, ($p < 10^{-4}$) as well as age over 50 (0.002), were independently associated with a cardiovascular emergency.

Table II: Crude and adjusted associations between patients' characteristics and cardiovascular emergencies in the 422 participants

Characteristics	Crude OR [95% CI]		Adjusted, ORa [95% CI]		p-value
Age, years					
<50	1		1		
≥50	2.74 [1.74-4.30]	<10 ⁻⁴	2.34 [1.33-4.10]		0.002
Personal history of hypertension					
No	1		1		
Yes	6.30 [3.87-10.25]	<10 ⁻⁴	6.09 [3.31-11.21]		<10 ⁻⁴
Body mass index, kg/m²					
< 30	1		1		
≥ 30	1.12 [0.60-2.11]	0.7			
Sex					
Male	1		1		
Female	0.82 [0.53-1.27]	0.38			
Physical activity practice					
No	1		1		
Yes	1.70 [1.09-2.64]	0.01			
Daily time spent sitting					
≤ 6h	1		1		
> 6h	1.38 [0.90-2.12]	0.13	1.56 [0.89-2.73]		0.11
Waist circumference[†], n=312					
≤ standard / sex	1		1		
> standard / sex	1.74 [1.06-2.84]	0.02	1.51 [0.86-2.67]		0.14
Use of oral contraceptives					
No	1		1		
Yes	0.55 [0.20-1.49]	0.23			
Smoking					
No	1		1		
Yes	0.59 [0.26-1.32]	0.19			
Spouse smoking, n=298					
No	1		1		
Yes	6.66 [1.27-34.86]	<10 ⁻⁴			
Diabetes					
No	1		1		
Yes	0.53 [0.22-1.23]	0.13	0.27 [0.09-0.76]		0.01

Abbreviations: OR: Odds ratio; ORa: Adjusted Odds ratio; CI: 95% confidence interval; †: for men, we considered a threshold below the standards as a waist circumference of <102 cm and < 88 for women cm.

DISCUSSION

We found a prevalence of cardiovascular emergencies of 27.5%. This estimate is higher than that found in the Teaching hospital Gabriel Touré (1.2%) by Diop et al. in Mali [9], and in hospitals in North Cameroon (9.9%) by Olivier Pancha et al. [10]. In one hand, their emergency departments even received surgical emergencies and the sampling was also exhaustive during a year. On the other hand, it is lower than the prevalence reported by Mboliasa et al. in Kinshasa which was 32.5% [11]. This latest study was carried out in an intensive care unit, which has only 8 beds, and expected to admit more life-threatening pathologies, rather than uncomplicated infectious ones. Doing so, that could overestimate the magnitude of cardiovascular emergencies. All these figures confirm the data of WHO which noted a clear progression of cardiovascular diseases in the world, this due to the epidemiological transition and the insufficiency or lack of preventive measures. In our study, the most common cardiovascular emergencies were: stroke: 37 patients (32%), hypertensive flare: 33 patients (28%) and heart failure: 21 patients (18%). The importance of these three (3) pathologies is consistent with Sub-Saharan statistics on their distribution. Indeed, Bernard et al. [3],

estimated them to be 32.2% for severe hypertension, 27.5% for heart failure and 20.3% for stroke.

The main cardiovascular pathology in the emergency department of the Niamey National Hospital was stroke. It was the same in Togo [12] and Libreville [13]. On the other hand in predominant pathology was coronary syndrome according to Sarr et al., [14]. It must be said that in their context, carrying out additional examinations made the difference because the study was conducted in the cardiological intensive care unit, which has the essentials in terms of diagnosis.

The average age was around 58 years, which is close to average age reported by Mboliasa in Kinshasa [11] and Essola et al. in Libreville [13]. Our average age is lower than that found in Nigeria by Iloh et al [15].

In developing countries, the most affected population is mostly young adults [3,16]. Unlike in developed countries where cardiovascular emergencies are more frequently described in the elderly [17]. The male predominance corroborates the results of other studies [3,11] where men represented 53,4% and 61,6% respectively. The hypothesis would be that in women, oestrogen has a protective effect until menopause [18].

Cardiovascular risk factors at Niamey National Hospital were largely unknown until now. In our study, age over

50 and history of hypertension are the risk factors statistically associated with cardiovascular pathologies (respectively, $p = 0.02$, $p < 10^{-4}$). The latter is an established risk factor as reported by Qureshi et al. in the Framingham study, where hypertension was associated with increased cardiovascular events [19]. This result has also been reported by other authors in Germany [20], and in Japan [21].

Age is a known risk factor that gradually increases the incidence of complications from aortic, coronary and then carotid atheroma and heart failure. This risk becomes significant from the age of 50 for men and 60 for women [22].

Surprisingly, diabetes appears to be protective factor in the onset of a cardiovascular emergency in this study. We hypothesize it may be due to the fact that these patients are generally well followed and therefore enough aware of the potential complications of their disease. They could also benefit regularly from advice on healthy living and it is not excluded that their drug treatment confers this protection regarding cardiovascular emergencies. They will usually go for causes other than a cardiovascular emergency. There is a compelling case for the beneficial effects of therapeutic lifestyle modification on the prevalence of cardiovascular complications in type 2 diabetes [23]. Nevertheless, we believe that these results require confirmation by a larger study with a more appropriate study design such as a case-control study or a hospital cohort study.

Hypertension is the most frequent personal medical history because it was found in 50.8% of patients, which is in line with reports from other studies [3,9,11].

LIMITATIONS

Low socioeconomic status of 61.2% of patients hampered the performance of some additional test essential for diagnosis (brain scan, cardiac ultrasound). Thus, practitioners limited their diagnosis to the term "suspicion", which could have resulted in a classification bias.

CONCLUSION

Prevalence of cardiovascular emergencies was 27.5%. The most frequent cardiovascular emergencies were stroke, hypertensive surge, and, heart failure with respectively 32%, 28%, and 18% with a third of recorded deaths, associated with stroke ischemic. Cardiovascular emergency prevention strategies that target people aged ≥ 50 years, those with a history of hypertension, may help reduce the frequency of cardiovascular emergencies. In addition, a larger study could also shed light on the role of diabetes in the frequency of cardiovascular emergencies at Niamey National Hospital.

AVAILABILITY OF DATA AND MATERIAL

The data are available under request to the authors and the National Hospital of Niamey Staff at bafatoum@gmail.com with copy to harouna.bako@yahoo.fr

DISCLOSURE STATEMENT

The authors declared no conflict of interest.

All the authors read the manuscript and they consent to submit for publication.

FUNDINGS

Fatoumata received support from the Burkina Field Epidemiology and Laboratory Training Program. A part from that, the study does not benefit from any special funding.

AUTHOR'S CONTRIBUTION

FMI wrote the study protocol. JK supervised and validated the study protocol. FMI collected the data and performed the statistical analyses under supervision of Jean Kaboré. JK, PYK, ML, JBO and NM revised the manuscript and gave critical inputs.

ACKNOWLEDGMENTS

The authors would like to fulfil a duty of gratitude to medical staff who facilitated access to data collection and to those who gave any support during the study. Specially, authors are grateful to the Epidemic Surveillance and Response Directorate, to both staff of the medical emergency department and the National Hospital of Niamey.

REFERENCES

1. WHO. À propos des maladies cardiovasculaires [Internet]. 2018 [cité 1 févr 2020]. Disponible sur: https://www.who.int/cardiovascular_diseases/about_cvd/fr/
2. OMS. Maladies non transmissibles [Internet]. 2018 [cité 1 févr 2020]. Disponible sur: <https://www.who.int/fr/news-room/fact-sheets/detail/noncommunicable-diseases>
3. Bertrand E, Muna WFT, Diouf SM, Ekra A, Kane A, Kingue S, et al. Urgences cardiovasculaires en Afrique subsaharienne. Arch Mal Coeur Vaiss. déc 2006;99(12):1159-65.
4. Noubiap JJ, Bigna J, Nansseu J, Nyaga U, Balti E, Echouffo-Tcheugui JB, et al. Prevalence of dyslipidaemia among adults in Africa: a systematic review and meta-analysis. Lancet Glob Health. 1 sept 2018;6(9):998-1007.
5. OMS. Maladies cardiovasculaires [Internet]. 2017 [cité 1 févr 2020]. Disponible sur: [https://www.who.int/fr/news-room/fact-sheets/detail/cardiovascular-diseases-\(cvds\)](https://www.who.int/fr/news-room/fact-sheets/detail/cardiovascular-diseases-(cvds))
6. Noubiap JJ ME. Prevalence of elevated blood pressure in children and adolescents in Africa: a systematic review and meta-analysis. Lancet Public Health. août 2017;375-86.
7. Adeloje D., Bigna J JR. An Estimate of the Incidence and Prevalence of Stroke in Africa: A Systematic Review and Meta-Analysis. PLOS ONE. 26 juin 2014;9(6):e100724.
8. Ministère de la Santé Publique du Niger. Plan Stratégique PNLMT Niger [Internet]. 2012 [cité 1 févr 2020]. Disponible sur: https://www.iccp-portal.org/system/files/plans/NER_B3_Plan%20Strat%20C3%A9gigue%20PNLMT%20Niger.pdf
9. Diop Th M MM, Almeimoune A, Dembélé AS, Diango MD. Urgences Cardio-Vasculaires Au service D'accueil Des Urgences Du Chu Gabriel Toure. Mali Medical 2018 Tome Xxxiii N°3 [Internet]. 2018 [cité 6 mai 2020]; Disponible sur: <http://www.malimedical.org/2018/1c.pdf>
10. Olivier Pancha Mbouemboue JON Adonis Koono Koono, Falmata Maha, Samuel Kingue. Place et Profil Évolutif des Maladies Cardiovasculaires en Milieu Hospitalier Nord Camerounais: Le Cas de L'Hôpital Régional de Ngaoundere |

Pancha Mbouemboue | HEALTH SCIENCES AND DISEASES. Vol 16, No 1 (2015) [Internet]. 2015 [cité 2 févr 2020]; Disponible sur: <https://www.hsd-fmsb.org/index.php/hsd/article/view/470>

11. Mboliasa I et al. Profil épidémiologique et clinique des urgences cardiovasculaires admises aux soins intensifs de médecine interne des Cliniques Universitaires de Kinshasa. *Ann Afr Médecine* [Internet]. mars 2015 [cité 1 févr 2020]; Volume 8 N°2. Disponible sur: <https://anafrimed.net/profil-epidemiologique-et-clinique-des-urgences-cardiovasculaires-admises-aux-soins-intensifs-de-medecine-interne-des-cliniques-universitaires-de-kinshasa/>

12. Damorou F PS Lawson B, Abdoulaye S, Soussou B, Grunitzky K. Urgences Cardio Vasculaires Et Leur Morbi mortalité En Milieu Hospitalier. A Propos De 733 Cas Colligés Au Chu Campus De Lomé (Hôpital De Reference National Du Togo) [Internet]. *Mali Médical* 2008 Tome Xxiii N°2. 2008 [cité 6 mai 2020]. Disponible sur: <http://malimedical.org/2008/55b.pdf>

13. Essola L MGM Ayo Bivigou. Les urgences cardiovasculaires au service d'accueil des urgences adultes du centre hospitalier universitaire de Libreville (chul). Volume 16 N° 44 : 63-69 [Internet]. 2018 [cité 1 févr 2020]; Disponible sur: <http://www.lebmo.org/les-urgences-cardiovasculaires-au-service-d%CA%BCaccueil-des-urgences-adultes-du-centre-hospitalier-universitaire-de-libreville-chul/>

14. Simon Antoine Sarr KB NanAjoua Archich, Bodian, MouHAMADou Bamba Ndiaye,, Kana. Profil des patients admis en unité de soins intensifs dans un service de cardiologie de Dakar. *Rev Afr Malgache Rech Sci Santé* [Internet]. 4 oct 2017 [cité 2 févr 2020];5(1). Disponible sur: <http://publication.lecames.org/index.php/sante/article/view/1126>

15. Iloh GUP, Amadi COA, Amadi AN. Epidemiology of Cardiovascular Emergencies in the Adult Emergency Department of a Tertiary Hospital in a Resource-constrained Setting of South-Eastern Nigeria. *J Adv Med Med Res.* 2 avr 2018;25(12): 1-10, 2018; Article no.JAMMR.25928:1-10.

16. Ellenga Mbolla Bf K-KE Kouala Landa Cm, Mongo Ngamami Sf, Makani Bassakouahou J, Ikama Ms, Ondze-Kafata I, Bakekolo Rp, Gombet Tra, Kimbally Kaky Sg. Les urgences cardiovasculaires dans le service de cardiologie et médecine interne du Centre Hospitalier et Universitaire de Brazzaville (Congo). *Cardiol Trop* N°143 Jan-Fev-Mars 2016 [Internet]. 2016 [cité 1 févr 2020]; Disponible sur: <http://tropical-cardiology.com/Accueil/index.php/2013-08-10-06-44-55/volume-n-143-livret-d-abstract/81-c14-les-urgences-cardiovasculaires-dans-le-service-de-cardiologie-et-medecine-interne-du-centre-hospitalier-et-universitaire-de-brazzaville-congo>

17. Cheval B RJ Eydoux N, et al. Prévalence des pathologies cardiovasculaires en médecine d'urgence dans le réseau Rescuer : données liées au sexe, à l'âge et au type de prise en charge. *Journal Européen des Urgences.* 2009;Volume-22, n° S2 page A96 (juin 2009).

18. Stanhewicz AE, Wenner MM, Stachenfeld NS. Sex differences in endothelial function important to vascular health and overall cardiovascular disease risk across the lifespan. *Am J Physiol Heart Circ Physiol.* 1 déc 2018;315(6):H1569-88.

19. Adnan I. Qureshi MFKS Jawad F Kirmani, Afshin A Divani, and Yousef Mohammad. Is Prehypertension a Risk Factor for Cardiovascular Diseases? | Stroke? *Stroke* 2005; :36:1859-1863.

20. P Trenkwalder PH R Schöniger, J Rossberg, H Lydtin, HW Hense. Hypertension as a risk factor for cardiovascular morbidity and mortality in an elderly German population. *European Heart Journal*, Volume 20, Issue 23, 1 December 1999, Pages 1752–1756, [Internet]. 1 déc 1999 [cité 7 juill 2020]; Disponible sur: <https://academic.oup.com/eurheartj/article/20/23/1752/481451>

21. Xiang M, Hu H, Imai T, Nishihara A, Sasaki N, Ogasawara T, et al. Association between anthropometric indices of obesity and risk of cardiovascular disease in Japanese men. *J Occup Health.* 21 nov 2019;

22. Collège des Enseignants de Cardiologie et Maladies Vasculaires. Facteurs de risque cardiovasculaire et prévention. :11.

23. L.Schlienger. Modifications thérapeutiques du mode de vie et prévention cardiovasculaire chez les sujets diabétiques de type 2. *Médecine Mal Métaboliques.* 1 févr 2019;13(1):27-35.